

Soft and Hard Jets in QCD

I.M. Dremin¹

Lebedev Physical Institute, Moscow 119991, Russia

Abstract

Multiplicity of sets of soft jets with energies ranging in some interval is determined. The possible role of collective effects is discussed.

Keywords: gluon, jet, multiplicity

PACS: 12.38 Bx

The phenomenon of jet emission is well known in QCD and firmly established in experiment. The two-jet events in e^+e^- -annihilation provide us with unique possibility to measure jets with fixed energy equal to that of colliding particles. In all other cases we have to deal with sets of jets with different energies. In three-jet events and in any high- p_t process the produced jets are somehow distributed in their energies. E.g., energies of gluon jets in three-jet process change from some lower limit determined by the requirement to separate this process to any value defined by experimental choice. Therefore, one has to deal with sets of jets with energies ranging in some interval.

Multiplicity of jets at a given energy has been calculated in QCD, and results agree quite well with experiment (see the reviews in [1, 2, 3]). This can be done for sets of jets as well [4]. The proper weights for jets with different energies in the set are provided by the QCD equations. The collective effects due to strings pulled apart during the separation of jets and color screening are somehow accounted in QCD. They lead, e.g., to different suppression of multiplicities between $q\bar{q}$ and qg pairs of jets. By measuring the multiplicity of sets of soft jets, this effect can become more pronounced.

To simplify the presentation, I consider gluodynamics (see also [5]). If the probability to create n particles² in a jet is denoted as P_n , the generating function G is defined as

$$G(z, y) = \sum_{n=0}^{\infty} P_n(y)(1+z)^n, \quad (1)$$

where z is an auxiliary variable, $y = \ln(p\Theta/Q_0) = \ln(2Q/Q_0)$ is the evolution parameter, defining the energy scale, p is the initial momentum, Θ is the angle of the divergence of the jet (jet opening angle), assumed here to be fixed, Q is the jet virtuality, $Q_0 = \text{const}$.

The gluodynamics equation for the generating function is written as

$$dG/dy = \int_0^1 dx K(x) \gamma_0^2 [G(y + \ln x) G(y + \ln(1-x)) - G(y)], \quad (2)$$

where

$$\gamma_0^2 = \frac{6\alpha_S}{\pi}, \quad (3)$$

α_S is the coupling strength and the kernel $K(x)$ is

$$K(x) = \frac{1}{x} - (1-x)[2-x(1-x)]. \quad (4)$$

¹Email: dremin@lpi.ru

The equation for mean multiplicities follows from eq. (2):

$$\langle n(y) \rangle' = \int_0^1 dx \gamma_0^2 K(x) (\langle n(y + \ln x) \rangle + \langle n(y + \ln(1 - x)) \rangle - \langle n(y) \rangle). \quad (5)$$

As follows from eq. (5), the first two terms in the brackets correspond to mean multiplicities of two subjets, and their sum is larger than the third term denoting the mean multiplicity of the initial jet. Therefore, the integrand is positive. This does not contradict to the statement that for a given event the total multiplicity is a sum of multiplicities in the two subjets because the averages in eq. (5) are done at different energies.

The scaling property of the fixed coupling QCD [5] allows to look for the solution of the equation (5) with

$$\langle n \rangle \propto \exp(\gamma y) \quad (\gamma = \text{const}) \quad (6)$$

The anomalous dimension γ is determined from (5) as

$$\gamma = \gamma_0^2 \int_0^1 dx K(x) (x^\gamma + (1 - x)^\gamma - 1) = \gamma_0^2 M_1(1, \gamma), \quad (7)$$

where

$$M_1(z, \gamma) = \int_0^z dx K(x) (x^\gamma + (1 - x)^\gamma - 1). \quad (8)$$

For small enough γ and γ_0 one gets

$$\gamma \approx \gamma_0(1 - 0.458\gamma_0 + 0.213\gamma_0^2). \quad (9)$$

Now, according to the above discussion we define soft jets as those with sum of energies of belonging to them particles less than some $x_0 E$. First, consider $x_0 = \text{const}$ and small. Then we should choose the upper limit of integration in eq. (5) equal to x_0 .

One gets from (5) the mean multiplicity of a set of soft jets $\langle n_s \rangle$:

$$\frac{\langle n_s \rangle}{\langle n \rangle} = \frac{M_1(x_0, \gamma)}{M_1(1, \gamma)}. \quad (10)$$

For small x_0 it is

$$\frac{\langle n_s \rangle}{\langle n \rangle} \approx \frac{\gamma_0^2}{\gamma^2} x_0^\gamma N_1(x_0, \gamma), \quad (11)$$

$$N_1(x_0, \gamma) = 1 - \gamma^2 x_0^{1-\gamma} - \frac{2\gamma}{1+\gamma} x_0 + \frac{\gamma^2(3+\gamma)}{4} x_0^{2-\gamma} + \frac{3\gamma}{2+\gamma} x_0^2 - \frac{\gamma^2(2+\gamma)}{3} x_0^{3-\gamma}. \quad (12)$$

Thus we have found the energy dependence of mean multiplicity of particles in a set of subjets with low energies $E_s \leq x_0 E$. As expected for constant x_0 , it is the same as the energy dependence of the total multiplicity with a different factor in front of it. Namely this dependence should be checked first in experimental data. Imposed on one another, these figures should coincide up to a normalization factor (10). This would confirm universality of gluons in jets.

Quite interesting is the non-trivial dependence of the normalization factor in eq. (10) on the parameter x_0 , which does not coincide simply with x_0^γ . It reflects the structure of QCD kernel $K(x)$. The main dependence on the cut-off parameter x_0 is given for $x_0 \ll 1$ by the factor x_0^γ with the same power as in dependence of total multiplicity on energy. This corresponds

eq. (12). They are induced by subjects with energies lower than $x_0 E$. The decrease of the normalization factor corresponds to diminishing role of very low energy jets at higher initial energies. This should be also checked in experiment.

If plotted as a function of the maximum energy in a set of jets ϵ_m , the mean multiplicity is

$$\langle n_s \rangle \propto \epsilon_m^\gamma \left[1 - \gamma^2 \left(\frac{\epsilon_m}{E} \right)^{1-\gamma} - \frac{2\gamma}{1+\gamma} \left(\frac{\epsilon_m}{E} \right) + \frac{\gamma^2(3+\gamma)}{4} \left(\frac{\epsilon_m}{E} \right)^{2-\gamma} + \frac{3\gamma}{2+\gamma} \left(\frac{\epsilon_m}{E} \right)^2 - \frac{\gamma^2(2+\gamma)}{3} \left(\frac{\epsilon_m}{E} \right)^{3-\gamma} \right]. \quad (13)$$

It reminds eq. (6) with the correction factor in the brackets.

This is the consequence of the scaling property of the fixed coupling QCD which results in the jets selfsimilarity.

In principle, other definitions of soft jets are possible with $x_0 = x_0(E)$. Then one should solve the equation

$$\frac{d\langle n_s \rangle}{dE} = E^{\gamma-1} \gamma_0^2 M_1(x_0(E), \gamma), \quad (14)$$

which follows from eq. (5). For example, one can choose the jets with energies less than some fixed constant independent of the initial energy. This would imply $\epsilon_m = \text{const}$ or $x_0(E) \propto 1/E$, and the exact integration of eq. (14) is necessary. However, for qualitative estimates, eqs (11)-(13) can be used. They show that the ratio of average multiplicities (11) tends to a constant at high energies corresponding to the multiplicity at the upper limit. At lower energies, it slightly increases with energy due to increasing role of jets with energies closest to their upper limit.

It is well known that for running coupling the power dependence $s^{\gamma/2}$ is replaced by $\exp(c\sqrt{\ln s})$. The qualitative statement about the similar energy behaviour of mean multiplicities in soft and inclusive processes should be valid also.

The above results can be confronted to experimental data if soft jets are separated in 3-jet events or in high- p_t hadronic collisions. However, in our treatment we did not consider the common experimental cut-off which must be also taken into account. This is the low-energy cut-off imposed on a soft jet for it to be observable. It requires the soft jet not to be extremely soft. Otherwise the third jet is not separated and the whole event is considered as a 2-jet one. Thus the share of energy must be larger than some x_1 , and the integration in eq. (8) should be from x_1 to x_0 . For $x_1 \leq x_0 \ll 1$ one gets

$$\frac{\langle n_s \rangle}{\langle n \rangle} = \frac{\gamma_0^2}{\gamma^2} [v(x_0) - v(x_1)], \quad (15)$$

where the function $v(x)$ is easily guessed from eqs (11), (12). At $x_1 \ll x_0 \ll 1$ eq. (11) is restored.

The values for hard jets are obtained by subtracting these results from values for the total process.

I concentrated here on mean multiplicity but similar calculations have been done [4] for higher moments of multiplicity distributions, and they can be compared with experiment.

The main problem in comparison is related to the jet definition, i.e., to treatment of particles belonging to the regions between the jets. They are more influenced by the collective effects which appear even in e^+e^- annihilation due to strings pulled between jets and their mutual color screening. The experimental verification of the normalization factor in (10) (for small x_0 it is the behavior of N_1 (11)) becomes important because it would show that these collective

Another interesting experimental aspect of collective effects is the energy distribution among jets in three-jet events. If boldly treated, the QCD diagram of the process implies that one of the quark jets remains untouched and must have the same energy as in two-jet events. Another (anti)quark jet emits a gluon, and they share this energy. Therefore, the energy distribution should have the two-bump structure with one bump at the primary energy and another one at smaller values (near half of it if the energy is shared equally between quarks and gluons). This statement is correct if collective effects due to string tension play minor role. In principle, string tension can lead to some collective energy flow from one jet to another one and change the shape of the energy distribution among three produced jets. Thus one can determine the role of collective effects by measuring this distribution.

In conclusion, the experimentally measured values of mean multiplicities of particles belonging to a set of soft jets can be compared with the obtained above theoretical predictions at different values of this share of energy. For a constant share, this dependence is the same as for the average total multiplicity but with non-trivial x_0 -dependence of the factor in front of it. Some predictions are obtained for energy dependent cut-offs. The collective effects due to string tension are discussed. These results can be confronted to experiment. The results concerning the behavior of higher moments analogous to those for fixed energy jets [6] will be published elsewhere.

Acknowledgments

This work has been supported in part by the RFBR grants 03-02-16134, 04-02-16445, NSH-1936.2003.2.

References

- [1] I.M. Dremin, UFN 164 (1994) 785; Physics-Uspekhi 37 (1994) 715.
- [2] I.M. Dremin and J.W. Gary, Phys. Rep. 349 (2001) 301.
- [3] V.A.Khoze and W. Ochs, Int. J. Mod. Phys. A 12 (1997) 2949.
- [4] I.M. Dremin, JETP Lett. 81 (2005) 391.
- [5] I.M. Dremin and R.C. Hwa, Phys. Lett. B 324 (1994) 477; Phys. Rev. D 49 (1994) 5805.
- [6] I.M. Dremin, V.A. Nechitailo, Mod. Phys. Lett. A 9 (1994) 1471; JETP Lett. 58 (1993) 881.